

VACCINATION CHART

AGE	NAME OF THE VACCINE	DETAILS
BIRTH	BCG	Administer BCG, OPV-O, HepB1 vaccine to all newborns before hospital discharge
	OPV O	
	Hep – B1	
6 WEEKS	IPV 1 / OPV	Polio: <ul style="list-style-type: none"> OPV can be given to children till 5 years of age. IPV is an injectable vaccine. Rotavirus: 2 doses of RV -1 & 3 doses of RV – 5.
	DPT / DTaP 1	
	Hib 1	
	Hep – B2	
	Rotavirus 1	
	PCV 1	
10 WEEKS	IPV 2 / OPV	PCV : Is given to prevent pneumococcal infections. Total of four doses of this vaccine given, the first one being at six weeks.
	DPT /DTaP 2	
	Hib 2	
	Rotavirus 2	
	PCV 2	
14 WEEKS	IPV 3 /OPV	DPT/DTaP = Diphtheria, Pertusis, Tetanus Vaccine Hib = Hemophilus Influenza B Vaccine Rotavirus: Only 2 doses of RV 1 are recommended at present.
	DPT / DTaP3	
	Hib 3	
	Rotavirus 3	
	PCV 3	
6 MONTHS	OPV 1	Hepatitis B : The final (3 rd or 4 th) dose in the Hep B vaccine series should be administered no earlier than age 24 weeks & atleast 16 weeks after the first dose.
	Hep – B3	
	Flu Vaccine 1	
	Flu Vaccine 2	
	Booster	
Flu vaccine booster Every Year		
9 MONTHS	OPV 2	An interval of atleast 4 weeks with the MMR vaccine should be maintained while administering Typhoid vaccine.
	MMR-1	
9 to 12 MONTHS	Typhoid Conjugate Vaccine	
12 MONTHS	Hep – A1	Hepatitis A: For both killed & live hepatitis A vaccines, 2 doses are recommended.
15 MONTHS	MMR -2	Varicella: The risk of breakthrough varicella is lower if given 15 months onwards.
	Varicella 1	
	PCV booster	
16 to 18 MONTHS	IPV B 1	The 1 st booster (4 th dose) may be administered as early as age 12 months, provided atleast 6 months have elapsed since the third dose.
	DTwP / DTaP 1 Booster	
	Hib B 1	
18 MONTHS	Hep – A2	Hepatitis A: For both killed & live hepatitis A vaccines, 2 doses are recommended.
2 YEARS	Typhoid Booster	
4 to 6 YEARS	DTwP B 2 / DTaP B 2	Tdap vaccine can protect adolescents and adults from tetanus, diphtheria, and pertussis. Tdap vaccine is also given to pregnant women. HPV: 2-dose schedule for adolescents who start the vaccination series before the age 15. The 2 doses should be separated by 6 to 12 months.
	OPV 3	
	Varicella 2	
	Typhoid Booster	
10 to 12 YEARS	Tdap / Td	
	HPV	